arXiv:1606.03706v2 [math.GT] 12 Oct 2016

VEECH GROUPS OF FLAT SURFACES WITH POLES

GUILLAUME TAHAR

Abstract. Flat surfaces that correspond to meromorphic 1-forms with poles or to meromorphic quadratic differentials containing poles of order two and higher have infinite flat area. We classify groups that appear as Veech groups of translation surfaces with poles. We characterize those surfaces such that their $GL^+(2,\mathbb{R})$-orbit or their $SL(2,\mathbb{R})$-orbit is closed. Finally, we provide a way to determine the Veech group for a typical infinite surface in any given chamber of a stratum.

Contents

1. Introduction 1
2. Statement of main results 2
3. Definitions and tools 4
4. Classification of Veech Groups 8
5. Closedness of orbits 10
6. Generic Veech groups 13
References 15

1. Introduction

Moduli spaces of translation structures on surfaces are endowed with an action of $GL^+(2,\mathbb{R})$. The Veech group of a flat surface X is the stabilizer of X by this group action. Characterization of Veech groups of translation surfaces corresponding to Abelian differentials is an open problem [3, 10, 15]. We show in this paper that flat surfaces corresponding to meromorphic differentials are rigid enough to allow a full solution to this problem.
Some results about the Veech group of a flat surface of infinite area are already known. In [13], Valdez computes the Veech group of some families of surfaces of infinite area associated to irrational billards. In [7], Hubert and Schmithüsen provide examples of square tiled surfaces with an infinite number of tiles and whose Veech groups are infinitely generated subgroups of $SL(2,\mathbb{R})$. In this paper, we consider only surfaces of finite type, that is surfaces of finite genus and whose singularities have a finite order.

Meromorphic 1-forms and quadratic differentials with poles of higher order induce flat structures of infinite area. However, they have finite complexity, see [1, 4] and we have some tools to study them [1, 12]. Since we cannot normalize the area, the action of $GL^+(2,\mathbb{R})$ does not reduce to that of $SL(2,\mathbb{R})$.

Let X be a compact Riemann Surface and q be a meromorphic quadratic differential with at least one pole of higher order, that is a pole of order no smaller than two. Such a pair (X,q) is called a *flat surface with poles of higher order*. When there is no ambiguity,

Date: October 12, 2016.
Key words and phrases. Translation surface, Veech Group, Flat structure.

it will be referred to as X. The meromorphic quadratic differential q is allowed to be the global square of a meromorphic differential so our study includes meromorphic 1-forms as a special case.

Zeroes and poles of order one of the quadratic differential are conical singularities for the induced flat metric [11]. We denote by $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$ the stratum in moduli space of meromorphic quadratic differentials that corresponds to meromorphic differentials with conical singularites of degrees $a_1, \ldots, a_n \in \{-1\} \cup \mathbb{N}^*$ and poles of degrees $b_1, \ldots, b_p \geq 2$. We have $\sum_{i=1}^{n} a_i - \sum_{j=1}^{p} b_j = 4g - 4$ where g is the *genus* of the underlying Riemann surface. Unless otherwise indicated, we will assume that $n > 0$ and $p > 0$. Besides, if $g = 0$, we will assume that $n + p \geq 3$. In order to simplify our study, there are no marked points.
In this paper, we consider the Veech group of a flat surface with poles of higher order (X, q) as a subgroup of $GL^+(2, \mathbb{R})/\{\pm Id\}$. When the quadratic differential is the global square of a meromorphic 1-form ω, what we get is a finite index subgroup of the Veech group of (X, ω) using the usual definition.

2. Statement of main results

There is a fundamental geometric invariant of translation surfaces with poles and their quadratic counterparts. The convex hull $core(X)$ of the set of conical singularities of a surface X is a kind of polygon, see [4, 12], whose boundary is a finite chain of saddle connections. The core is equivariant for the action of the Veech group. Therefore, symmetries of the surface are significantly restricted. For instance, in most cases, the $GL^+(2, \mathbb{R})$ action is not ergodic [2].
Besides, unlike strata of translation surfaces, strata of flat surfaces with poles of higher order have a walls-and-chambers structure defined by topological changes of the core. The cores of two surfaces that belong to the same chamber are topologically identical.

Our main theorem is a complete classification of Veech groups of flat surfaces with poles of higher order. Valdez proved an analogous classification in the context of Abelian differentials on stable curves, see [13].

Theorem 2.1. *The Veech group of a flat surface with poles of higher order belongs to one of the following three types of subgroups of $GL^+(2, \mathbb{R})/\{\pm Id\}$:*
- Finite type: conjugated to a finite rotation group ;
- Cyclic parabolic type: conjugated to $\left\{ \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} \mid k \in \mathbb{Z} \right\}$;
- Continuous type: conjugated to $\left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a \in \mathbb{R}_+^*, b \in \mathbb{R} \right\}$.

Theorem 2.1 is proved in section 4.

For classical translation surfaces, the $GL^+(2, \mathbb{R})$-orbit is closed if and only if the Veech group is a lattice subgroup in $SL(2, \mathbb{R})$, see [14]. There is an analogous result for translation surfaces with poles.

Theorem 2.2. *The following statements are equivalent for a flat surface with poles of higher order (X, q):*
(i) The Veech group of (X, q) is of continuous type.

(ii) All saddle connections of (X, q) share the same direction.
(iii) The $GL^+(2,\mathbb{R})$-orbit of (X, q) is closed in the ambient stratum.

As we cannot normalize the area of the surface, we have to distinguish the action of $GL^+(2,\mathbb{R})$ and that of $SL(2,\mathbb{R})$.
Surfaces whose $SL(2,\mathbb{R})$-orbit is closed are very specific among classical translation surfaces. They are called *Veech surfaces*. On the contrary, it is common for a translation surface with poles to have a closed $SL(2,\mathbb{R})$-orbit.

Theorem 2.3. *The $SL(2,\mathbb{R})$-orbit of a flat surface with poles of higher order (X, q) is closed in the ambient stratum in these two cases:*
(i) Not all saddle connections of $\partial\mathcal{C}(X)$ share the same direction.
(ii) All saddle connections of $\partial\mathcal{C}(X)$ share the same direction and $core(X)$ decomposes into a (maybe empty) family of cylinders of commensurable moduli.
Otherwise, the $SL(2,\mathbb{R})$-orbit of (X, q) is not closed in the ambient stratum.

Theorems 2.2 and 2.3 are proved in section 5.

Following definitions of [8], we say that a flat surface (X, q) is generic in its stratum (or its chamber) if it lies outside a countable union of real codimension one submanifolds. Two saddle connections are said to be parallel when their relative homology classes are not linearly independant over $\mathbb{R}$.

Generic Veech groups of flat surfaces of finite area are already known, see [8]. They are trivial as long as $g \geq 2$. We get similar results for flat surfaces with poles of higher order.

Theorem 2.4. *Let $\mathcal{C}$ be a chamber where the core is a unique finite volume cylinder in the following strata:*
- $\mathcal{Q}(a, b, -a-2, -b-2)$ with $a, b \geq 1$ (genus zero)
- $\mathcal{Q}(a, -1^2, -a-2)$ with $a \geq 1$ (genus zero)
- $\mathcal{Q}(a, -a)$ with $a \geq 2$ (genus one)
The Veech group of every flat surface with poles of higher order of $\mathcal{C}$ is of cyclic parabolic type.
In strata $\mathcal{Q}$ of surfaces of genus zero such that $n + p = 3$, the Veech group of every flat surface with poles of higher order is of continuous type.
Outside these chambers and strata, the Veech group of a generic surface is trivial.

Theorem 2.4 is proved in section 6.

The structure of the paper is the following:
- In Section 3, we recall the background and tools useful to study flat surfaces of infinite area : flat metric, saddle connections, the moduli space, the core of a surface and its associated wall-and-chambers structure, the contraction flow.
- In Section 4, we prove our theorem of classification.
- In Section 5, we provide complete criteria of closedness of orbits of $GL^+(2,\mathbb{R})$ and $SL(2,\mathbb{R})$.
- In Section 6, we characterize loci where the Veech group of a generic surface is nontrivial.

Acknowledgements. I thank my doctoral advisor Anton Zorich for motivating my work on this paper and many interesting discussions. I am grateful to Corentin Boissy, Quentin Gendron and Ben-Michael Kohli for their valuable remarks. This work is supported by the ERC Project "Quasiperiodic" of Artur Avila.

3. Definitions and tools

3.1. **Flat structures.** Let X be a compact Riemann surface and let q be a meromorphic quadratic differential. Λ and Δ respectively are the set of conical singularities and poles of higher order of q. Outside Λ and Δ, q is locally the square of a holomorphic differential ω. Integration of ω in a neighborhood of z_0 gives local coordinates whose transition maps are of the type $z \mapsto \pm z + c$. The pair (X, q) seen as a smooth surface with such a translation atlas is called a *flat surface with poles of higher order*.

In the case of quadratic differentials that are the global square of a 1-form, there is no holonomy of the metric. Otherwise, the holonomy of the metric is $\mathbb{Z}/2\mathbb{Z}$.

In a neighborhood of an element of Λ, the metric induced by q admits a conical singularity of angle $(k+2)\,\pi$ where k is the degree of the corresponding zero of q. In particular, poles of order one are conical singularities of angle π.

See Strebel [11] for a complete description of the local geometry around poles of higher order.

Definition 3.1. A saddle connection is a geodesic segment joining two conical singularities of the flat surface such that all interior points are not conical singularities.

In [12], we proved a lower bound on the number of saddle connections of a flat surface with poles of higher order.

Theorem 3.2. *Let $|SC|$ be the number of saddle connctions of a flat surface with poles of higher order (X, q) of genus g belonging to $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$, then we have $|SC| \geq 2g - 2 + n + p$.*

3.2. **Moduli space.** If (X, q) and (X', q') are flat surfaces such that there is a biholomorphism f from X to X' such that q is the pullback of q', then f is an isometry for the flat metrics defined by q and q'.

We define the moduli space of meromorphic quadratic differentials as the space of equivalence classes of pairs (X, q) up to biholomorphism preserving the quadratic form.

We denote by $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$ the *stratum* that corresponds to meromorphic quadratic forms with singularities of degrees $a_1, \ldots, a_n$ and $-b_1, \ldots, -b_p$. We have $a_1, \ldots, a_n \in \{-1\} \cup \mathbb{N}^*$. They are the poles of order one and the zeroes of arbitrary order. They define conical singularities. Singularities are referred to as poles of higher order when they are poles of order no smaller than two. That is why we have $b_1, \ldots, b_p \geq 2$.

3.3. **Canonical double covering.** Outside the set of singularities, every quadratic differential is locally the square of a 1-form. The canonical double covering assigns to every pair (X, q) of a Riemann surface and a quadratic differential a flat surface such that the quadratic differential is globally the square of a 1-form. For quadratic surfaces that are already the global square of a 1-form, instead of constructing a canonical double covering, we simply choose a square root.

Let (X, q) be a flat surface of $\mathcal{Q}(2c_1, \ldots, 2c_s, 2d_1 + 1, \ldots, 2d_t + 1)$ where $c_1, \ldots, c_s \in \mathbb{Z}^*$ and $d_1, \ldots, d_t \in \mathbb{Z}$. We denote by $P_i \in X$ the point corresponding to the pole of order

$2d_i + 1$.

We consider the double covering branched only over the set $\{P_1, \ldots, P_t\}$. The flat geometry of the resulting surface is that of the global square of a meromorphic differential ω. We denote by $(\widetilde{X}, \omega)$ the canonical double cover of (X, q). $(\widetilde{X}, \omega)$ belongs to $\mathcal{H}(c_1, \ldots, c_s, c_1, \ldots, c_s, 2d_1 + 2, \ldots, 2d_t + 2)$.

Quadratic differentials that are the global square of a 1-form belong to strata of the form $\mathcal{Q}(2c_1, \ldots, 2c_s)$ where all singularities are of even order. Their square roots belong to stratum $\mathcal{H}(c_1, \ldots, c_s)$.

3.4. **Period coordinates.** The double canonical covering $(\widetilde{X}, \omega)$ of (X, q) admits an involution τ that induces another involution τ^* on $H_1(X \setminus \Delta, \Lambda)$ where Δ is the set of poles of higher order and Λ is the set of conical singularities.
Thus $H_1(X \setminus \Delta, \Lambda)$ decomposes into an invariant subspace H_+ and an anti-invariant space H_-. Strata are complex-analytic orbifolds with local coordinates given by the period map of H_- [1].

We can associate homology classes to saddle connections. In the case of quadratic differentials, the way to do that is somewhat subtle. We consider a flat surface with poles (X, q) and its double canonical covering $(\widetilde{X}, \omega)$.

$H_1(\widetilde{X} \setminus \Delta', \Lambda')$ is the first relative homology group of $(\widetilde{X}, \omega)$ where Δ' and Λ' respectively are the preimages by π of the conical singularities and poles of higher order associated to q. An involution τ associated to the covering acts on $H_1(\widetilde{X} \setminus \Delta', \Lambda')$.

We denote by γ_1 and γ_2 the two preimages of an oriented saddle connection γ. If the relative cycle $[\gamma_1] \in H_1(\widetilde{X} \setminus \Delta', \Lambda')$ satisfies $[\gamma_1] = -[\gamma_2]$, then we define $[\gamma] = [\gamma_1] - [\gamma_2]$. Conversely, if $[\gamma_1] = [\gamma_2]$, then we define $[\gamma] = [\gamma_1] + [\gamma_2]$.
Two saddle connections are said to be parallel when their relative homology classes are linearly dependant over $\mathbb{R}$.
The holonomy vector of a saddle connection is essentially determined by the period of its relative homology class. Its direction modulo π is that of the period and its length is one-half of the modulus of the period.

3.5. **Parallel saddle connections.** We say that a couple γ_1, γ_2 of saddle connections is rigid when every deformation inside a neighborhood of the surface in the ambient stratum preserves the ratio $|\gamma_1| : |\gamma_2|$ between their lengths. The following lemma is proved as a part of Proposition 1 in [9]. Its proof generalizes without any difficulty in the case of meromorphic quadratic differentials with poles of arbitrary order.

Lemma 3.3. *Let (X, q) be a flat surface with (or without) poles of higher order, a couple γ_1, γ_2 of saddle connections is rigid if and only if γ_1 and γ_2 are parallel.*

Another lemma is needed to prove our results about generic Veech groups (Theorem 2.4 that is proved in Section 6). The following lemma is proved in [9] as Theorem 1. Like in the previous case, its proof generalizes straightforwardly in the case of meromorphic quadratic differentials with poles of arbitrary order.

Lemma 3.4. *Let (X, q) be a flat surface with (or without) poles of higher order. Two saddle connections γ_1 and γ_2 are parallel if and only if they have no interior intersections*

and one of the connected components of $X \setminus (\gamma_1 \cup \gamma_2)$ *has trivial linear holonomy.*

3.6. **Index of a loop.** The topological index of a loop is particularly easy to handle in flat geometry.

Definition 3.5. Let γ be a simple closed curve in a flat surface with (or without) poles of higher order. γ is parametrized by arc-length and passes only through regular points. We consider the lifting η of γ by the canonical double covering. Then $\eta'(t) = e^{i\theta(t)}$.
We have $\frac{1}{2\pi}\int_0^T \theta'(t)dt \in \frac{1}{2}\mathbb{Z}$ because of the holonomy of the flat surface. This number is the topological index $ind(\gamma)$ of our loop γ.

In particular, the topological index of a simple closed curve around a singularity of order k is $1 + \frac{k}{2}$.

3.7. **Core of a flat surface with poles of higher order.** The core of a flat surface of infinite area was introduced in [4] and systematically used in [12]. It is the convex hull of the conical singularities of a flat surface. Since all saddle connections belong to it, the core encompasses most of the qualitative (see walls-and chambers structure in subsection 3.7) and quantitative (periods of the homology) information about the geometry of the flat surface.

Definition 3.6. A subset E of a flat surface is *convex* if and only if every element of any geodesic segment between two points of E belongs to E.
The core of a flat surface with poles of higher order (X, q) is the convex hull $core(X)$ of its conical singularities Λ.
$\mathcal{IC}(X)$ is the interior of $core(X)$ in X and $\partial\mathcal{C}(X) = core(X) \setminus \mathcal{IC}(X)$ is its boundary.
The *core* is said to be degenerated when $\mathcal{IC}(X) = \emptyset$ that is when $core(X)$ is just graph $\partial\mathcal{C}(X)$.

Lemma 3.7. *Let* X *be a flat surface with poles of higher order* $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$*, then* $X \setminus core(X)$ *has* p *connected components. Each of them is a topological disk. We refer to these connected components as domains of poles.*

Proof. Following Proposition 2.3 in [4], $core(X)$ is a deformation retract of $X \setminus \Delta$ where Δ is the set of poles of higher order. □

Lemma 3.8. *For any flat surface with poles of higher order* X*,* $\partial\mathcal{C}(X)$ *is a finite union of saddle connections.*

Proof. See Proposition 2.2 in [4].

□

3.8. **Discriminant and walls-and-chambers structure.** Strata of flat surfaces with poles of higher order decompose into chambers where the qualitative shape of the core is the same. The discriminant is the locus that separates these chambers from each other.

Definition 3.9. A flat surface with poles of higher order X belongs to the discriminant of the stratum if and only if two nonparallel consecutive saddle connections of the boundary of the core share an angle of π. Chambers are defined to be the connected components of the complementary to the discriminant in the strata.

The following lemma is proved as Proposition 4.12 in [12].

Lemma 3.10. *The discriminant is a $GL^+(2,\mathbb{R})$-invariant hypersurface of real codimension one in the stratum.*

The topological map on a flat surface with poles of higher order (X, q) defined by the embedded graph $\partial\mathcal{C}(X)$ is invariant along the chambers. The qualitative shape of the core and in particular the number of saddle connections of its boundary depend only on the chamber (see Proposition 4.13 in [12] for details).

3.9. **Dynamics and decomposition into invariant components.** We essentially follow the definitions Strebel gives in [11].

Definition 3.11. Depending on the direction, a trajectory starting from a regular point is of one of the four following types:
- regular closed geodesic (the trajectory is periodic),
- critical trajectory (the trajectory reaches a zero in finite time),
- trajectory finishing at a pole (the trajectory converges to a pole as $t \to +\infty$),
- recurrent trajectory (infinite trajectory nonconverging to a pole).

Theorem 3.12 describes how the directional flow decomposes flat surfaces into a finite number of invariant components.

Theorem 3.12. *Let X be a flat surface of genus g with poles of higher order in a stratum $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$. Cutting along all saddle connections sharing a given direction θ, we obtain finitely many connected components called invariant components. There are four types of invariant components:*
- ***finite volume cylinders*** *where the leaves are periodic with the same period,*
- ***minimal components*** *of finite volume where the foliation is minimal, the directions are recurrent and whose dynamics are given by a nontrivial interval exchange map,*
- ***infinite volume cylinders*** *bounding a simple pole and where the leaves are periodic with the same period,*
- ***free components*** *of infinite volume where generic leaves go from a pole to another or return to the same pole.*

Finite volume components belong to core(X). The number of finite volume invariant components in a given direction is bounded by $g + n - 1$.

Proof. Theorem 2.3 in [12] proves most of the theorem. Theorem 2.13 in [12] provides the bound on the number of finite volume invariant components.

□

3.10. $GL^+(2,\mathbb{R})$ **action and contraction flow.** On each stratum, $GL^+(2,\mathbb{R})$ acts by composition with coordinate functions, see [15]. Since neighborhoods of poles of higher order have infinite area, we cannot normalize the area of the surface and thus must consider the full action of $GL^+(2,\mathbb{R})$.

For a flat surface (X, q), the *stabilizer* $stab(X) \subset GL^+(2,\mathbb{R})$ is the subgroup of those $g \in GL^+(2,\mathbb{R})$ for which $gX = X$. The quotient $stab(X)/\{\pm Id\} \subset GL^+(2,\mathbb{R})/\{\pm Id\}$ is called the *Veech Group* of the flat surface (X, q).
Veech groups of two surfaces belonging to the same $GL^+(2,\mathbb{R})$-orbit are conjugated.

The contraction flow is a tool that allows to construct surfaces with degenerated core in a systematic way. All foundation results are already proved in [12].

Definition 3.13. Let α and θ be two distinct directions. The contraction flow is the action of the semigroup of matrices $C^t_{\alpha,\theta}$ conjugated to $\begin{pmatrix} e^{-t} & 0 \\ 0 & 1 \end{pmatrix}$ such that α is the contracted direction and θ is the preserved direction.

Lemma 3.14. *Let (X, q) be a flat surface with poles of higher order. θ is a direction that is not a direction of a saddle connection and α is any direction different from θ. The sequence $C^t_{\alpha,\theta}(X, q)$ converges to a surface (X_0, q_0) in the ambiant stratum. All saddle connections of (X_0, q_0) share the same direction θ and $core(X_0)$ is degenerated.*

Proof. See Lemma 2.2 of [12].

□

4. Classification of Veech Groups

Veech groups of flat surfaces with poles of higher order are very different from those of usual translation surfaces. They are either very big (not discrete) or very small (virtually cyclic). For instance, they cannot be lattices so there are no Veech surfaces with poles.

Proof of Theorem 2.1. Let (X, q) be a flat surface with poles of higher order. There is a finite number of saddle connections in $\partial\mathcal{C}(X)$. The holonomy vectors of these saddle connections are a finite subset of $\mathbb{C}/\{\pm 1\}$. Since this set is preserved by the action of the Veech group, either all saddle connections of $\partial\mathcal{C}(X)$ share the same direction or the Veech group is a finite group. Finite subgroups of $GL^+(2, \mathbb{R})$ are rotation groups. We say in this case that the Veech group is of finite type.

In the following, we consider that all saddle connections of $\partial\mathcal{C}(X)$ share the same direction θ. This direction is unchanged by the action of the Veech group and the lengths of these saddle connections are preserved. Therefore, in this case, the Veech group is conjugated to a subgroup of $\left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbb{R} \right\}$.

The next disjonction is between surfaces with degenerated and nondegenerated core. If $core(X)$ is degenerated, then every saddle connection belongs to $\partial\mathcal{C}(X)$ and by hypothesis they belong to the same direction. Without loss of generality, we can consider they all belong to the vertical direction. The whole geometry of X is that of infinite horizontal strips whose intersection with $core(X)$ are vertical saddle connections. Thus, the action of every element of $GL^+(2, \mathbb{R})$ preserving the vertical direction leaves (X, q) unchanged. Thus, the Veech group of (X, q) is conjugated to $\left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbb{R} \right\}$.

If $core(X)$ is not degenerated, then there are two saddle connections that have different directions. The set of holonomy vectors of the saddle connections of the whole surface is a discrete subset S of $\mathbb{C}/\{\pm 1\}$ that is preserved by the action of the Veech group. As there are two saddle connections that have different directions, then an element of the Veech group is entirely characterized by its action on S. Therefore, the Veech group of X is a discrete subgroup of $GL^+(2, \mathbb{R})/\{\pm Id\}$. This implies in particular that the Veech group of (X, q) is a discrete subgroup of $SL(2, \mathbb{R})/\{\pm Id\}$. We know it is also conjugated to a subgroup of $\left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbb{R} \right\}$. Therefore, either it is conjugated to $\left\{ \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} \mid k \in \mathbb{Z} \right\}$ or it is the trivial group.

□



FIGURE 1. A flat surface in $\mathcal{Q}(8, -2^6)$ with a discrete rotational symmetry.

One can find a realization for every Veech group of the classification provided by Theorem 2.1. The following example shows in particular that every cyclic finite group is realizable as a Veech group.

Example 4.1. Gluing infinite cylinders on the edges of a regular $2k$-gon, we get a surface in $\mathcal{Q}(4k-4, -2^{2k})$ whose Veech group is conjugated to $\mathbb{Z}/k\mathbb{Z}$, see Figure 1.

In the following, we characterize surfaces with Veech groups of cyclic parabolic type in terms of the decomposition in invariant components.

Proposition 4.2. *The Veech group of a flat surface with poles of higher order X is of cyclic parabolic type if and only if $core(X)$ is not degenerated and admits a decomposition in a finite number of cylinders with commensurable moduli in some direction θ.*

Proof. For classical translation surfaces (and half-translation surfaces), if there is a parabolic element of the Veech group preserving a direction θ, then the surface decomposes into a finite number of cylinders parallel to θ and whose moduli are commensurable [5].
It is enough te be sure that for flat surfaces with poles of higher order, existence of a parabolic element of the Veech group preserving a direction θ implies there is no minimal component for direction θ.
Besides, the parabolic element preserves the boundary of $core(X)$. As it is a finite union of saddle connections, all of them share the same direction θ.
Then we consider a decomposition into components invariant by the flow in direction θ (Theorem 3.12). We get a finite number of finite volume cylinders and a finite number of infinite area invariant components. There is at least one cylinder because otherwise all saddle connections would be parallel and the Veech group would be of infinite type.
Since there is a parabolic element preserving direction θ, finite volume cylinders whose waist curves share direction θ have commensurable moduli. This is a well known result about translation surfaces (and half-translation surfaces), see Veech in [14].

Conversely if the boundary, the residues and commensurable cylinder share the same direction θ, then there is a parabolic element of $GL^+(2,\mathbb{R})$ that preserves $core(X)$. As an element preserving direction θ preserves domains of poles (whose boundary is a chain of saddle connections in direction θ), this parabolic element preserves the whole surface. Following Theorem 2.1 and assuming the core is not degenerated, this implies that the Veech group of the surface is of cyclic parabolic type. □



FIGURE 2. A flat surface in $\mathcal{Q}(8, -2^4)$ whose Veech group is of cyclic parabolic type.



FIGURE 3. A surface in $\mathcal{Q}(4, -2^2)$ and the surface obtained through the use of the contraction flow along a given direction.

Example 4.3. We can construct many surfaces with a Veech group of cyclic parabolic type starting from a square-tiled core, see Figure 2.

Starting from any surface, the contraction flow in a generic direction provides examples of surfaces with a Veech group of infinite type, see Figure 3.

The contraction flow provides many examples of surfaces with a Veech group of infinite type. However, many of these surfaces belong to the discriminant of the ambiant stratum. The following example shows that this is not always true.

Example 4.4. Three flat planes with a vertical slit in each of them and connected cyclically provide a surface in $\mathcal{Q}(4^2, -4^3)$ with a Veech group of continuous type and that does not belong to the discriminant.

5. Closedness of orbits

In the study of classical translation surfaces, the Veech surfaces have a particular importance because they are the class of surfaces such that their $GL^+(2, \mathbb{R})$-orbit is closed, see [6]. Here, we characterize the class of flat surfaces with poles of higher order that share such a property. Theorem 2.2 proves that surfaces whose $GL^+(2, \mathbb{R})$-orbit is closed are exactly the surfaces whose Veech group is of continuous type.

Proof of Theorem 2.2. We first prove that (i) implies (iii). Let G be the Veech group of continuous type of a surface (X,q). $\mathbb{C}^* \cong \left\{ \begin{pmatrix} acos(\theta) & -asin(\theta) \\ asin(\theta) & acos(\theta) \end{pmatrix} \mid a \in \mathbb{R}^*,\ \theta \in \mathbb{R} \right\}$ and G generate $GL^+(2,\mathbb{R})$. So the action of $\mathbb{C}^*$ by rotations and homotheties is an affine parametrization of $GL^+(2,\mathbb{R}).(X,q)$. Therefore, $GL^+(2,\mathbb{R}).(X,q)$ is an affine plane that is closed.

Then we prove that (iii) implies (ii). If there are two saddle connections of different directions in a flat surface with poles of higher order (X,q), pushing the surface (X,q) through the contraction flow along a direction θ that is not a direction of saddle connection gives in the limit a surface (X_0,q_0) whose saddle connections share the same direction θ. So $(X_0,\omega_0) \notin GL^+(2,\mathbb{R}).(X,q)$ which is not closed. We proved by contraposition that (iii) implies (ii).

Finally, a flat surface (X,q) such that all saddle connections share the same direction has a Veech group that is of continuous type, see the proof of Theorem 2.1 for details. Therefore (ii) implies (i). This ends our circular reasoning.

□

While surfaces with a closed $GL^+(2,\mathbb{R})$-orbit are quite exceptional, most flat surfaces with poles of higher order have a closed $SL(2,\mathbb{R})$-orbit. Theorem 2.3 characterizes the surfaces whose $SL(2,\mathbb{R})$-orbit is closed in the ambient stratum.

Proof of Theorem 2.3. First, we prove that if a surface X belongs to the $SL(2,\mathbb{R})$-orbit closure of a surface X_0, then the topological types of $\partial\mathcal{C}(X)$ and $\partial\mathcal{C}(X_0)$ are the same. A change in the topological type of the core corresponds to a conical singularity that crosses a saddle connection belonging to the boundary of the core. Thus, it corresponds to degeneration of some ideal triangle. Since the action of $SL(2,\mathbb{R})$ is area-preserving, this cannot happen. Therefore, cores of all surfaces in a $SL(2,\mathbb{R})$-orbit closure share the same topological type.

We consider a flat surface (X,q) with poles of higher order such that there are two saddle connections of different directions in $\partial\mathcal{C}(X)$. As their holonomy is defined with an ambiguity of ± 1, we consider one of the representatives in $\mathbb{C}$ for each of these two saddle connections. We denote by w and z the representatives. They form a basis of $\mathbb{C}$ as a $\mathbb{R}$-vector space. Every surface (Y,q') in the $SL(2,\mathbb{R})$-orbit closure of (X,q) arises as the limit of a sequence $A_n.(X,q)$ where $\forall n \geq 0$, we have $A_n \in SL(2,\mathbb{R})$. As there are finitely many saddle connections in $\partial\mathcal{C}(Y)$, sequences $A_n.w$ and $A_n.z$ converge respectively to z_∞ and w_∞ in $\mathbb{C}$. Since matrices in $SL(2,\mathbb{R})$ are area-preserving, (z_∞, w_∞) is a basis of $\mathbb{C}$ as well. A matrix of $SL(2,\mathbb{R})$ is characterized by the image of a basis. Thus, sequence A_n converges to a limit $A_\infty \in SL(2,\mathbb{R})$. Therefore, $(Y,q') = A_\infty(X,q)$. In other words, every surface in the $SL(2,\mathbb{R})$-orbit closure of (X,q) actually belongs to the orbit of (X,q).

If the Veech group of a flat surface with poles of higher order (X,q) is of continuous type, then $GL^+(2,\mathbb{R}).(X,q)$ is closed (Theorem 2.3). As $SL(2,\mathbb{R})$ is a closed subgroup of $GL^+(2,\mathbb{R})$, $SL(2,\mathbb{R}).(X,q)$ is closed too. This covers situations where all saddle connections of $\partial\mathcal{C}(X)$ share the same direction and $core(X)$ is degenerated.

Then we consider surfaces (X,q) such that all saddle connections of $\partial\mathcal{C}(X)$ share the same direction θ and $core(X)$ is not degenerated. Without loss of generality, we suppose that θ is the vertical direction. We are going to use the Iwasawa decomposition of $SL(2,\mathbb{R})$. Every $M \in SL(2,\mathbb{R})$ has a unique representation as $M = K.A.N$ where we have

$K = \begin{pmatrix} cos(\alpha) & -sin(\alpha) \\ sin(\alpha) & cos(\alpha) \end{pmatrix}$, $A = \begin{pmatrix} r & 0 \\ 0 & 1/r \end{pmatrix}$ and $N = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$ with $\alpha, x \in \mathbb{R}$ and $r > 0$. In particular, $K \in SO(2, \mathbb{R})$.

Every surface (Y, q') in the $SL(2, \mathbb{R})$-orbit closure of (X, q) arises as the limit of a sequence $M_n.(X, q)$ where $\forall n \geq 0$, we have $M_n \in SL(2, \mathbb{R})$. Besides, $\forall n \geq 0$, we have $M_n = K_n A_n N_n$ in the Iwasawa decomposition.

First, $SO(2, \mathbb{R})$ is compact so there is $\alpha_\infty \in \mathbb{R}$ and an increasing subsequence $(\phi(n))_{n\geq 0}$ such that we have:

$$\lim_{n\to+\infty} K_{\phi(n)} = \begin{pmatrix} cos(\alpha_\infty) & -sin(\alpha_\infty) \\ sin(\alpha_\infty) & cos(\alpha_\infty) \end{pmatrix} \in SO(2, \mathbb{R}).$$

We have $A_n = \begin{pmatrix} r_n & 0 \\ 0 & 1/r_n \end{pmatrix}$. Since all saddle connections of $\partial\mathcal{C}(X)$ share the same direction, if there is an increasing subsequence $(\mu(n))_{n\geq 0}$ such that $\lim_{n\to+\infty} r_{\mu(n)} = +\infty$, then the length of the longest saddle connection in the boundary of the core of $M_{\mu(n)}.(X, q)$ grows to infinity. Therefore, $(r_n)_{n\geq 0}$ is bounded. For the same reasons (no saddle connection can shrink to saddle connection of zero length), $(r_n)_{n\geq 0}$ is bounded below by a positive number. That is why there is $r_\infty > 0$ and an increasing subsequence $(\psi(n))_{n\geq 0}$ such that $\lim_{n\to+\infty} r_{\psi\circ\phi(n)} = r_\infty$.
To recapitulate, we get an increasing subsequence $(\phi \circ \psi(n))_{n\geq 0}$ such that we have:

$$\lim_{n\to+\infty} K_{\phi\circ\psi(n)}.A_{\phi\circ\psi(n)} = \begin{pmatrix} cos(\alpha_\infty) & -sin(\alpha_\infty) \\ sin(\alpha_\infty) & cos(\alpha_\infty) \end{pmatrix} . \begin{pmatrix} r_\infty & 0 \\ 0 & 1/r_\infty \end{pmatrix}.$$

Therefore, the $SL(2, \mathbb{R})$-orbit of (X, q) is closed if and only if its H-orbit is closed where $H = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, x \in \mathbb{R} \right\}$.

If $core(X)$ decomposes into finite volume cylinders with commensurable moduli, then the Veech group of (X, q) is of cyclic parabolic type (Proposition 4.2) and the H-orbit of (X, q) is closed. Then, its $SL(2, \mathbb{R})$-orbit is closed too.

The last case is about surfaces (X, q) such that all saddle connections of $\partial\mathcal{C}(X)$ share the same direction (without loss of generality, we suppose this is the vertical direction) and $core(X)$ does not decompose into a family of cylinders of commensurable moduli. Either there is a minimal component or two cylinders whose moduli are not commensurable. Following Proposition 4.2, the Veech group of such a surface (X, q) is not of cyclic parabolic type and its H-orbit is not periodic.
Let $\mathcal{U}$ be the interior of $core(X)$. Since the boundary of $\mathcal{U}$ is made of vertical saddle connections whose total holonomy is zero, we can glue the boundary components on each other while adding marked points if necessary or modifying the order of the conical singularities. We get a flat surface (X_c, q_c) without poles of higher order but with some marked vertical saddle connections (those that have been glued on each other). Flat surface (X_c, q_c) belongs to a stratum $\mathcal{Q}$ of meromorphic quadratic differentials with at most simple poles.
No saddle connection shrinks under the action of H. Thus, lengths of saddle connections remain bounded below by a positive number all along the orbit. Therefore $H.(X_c, q_c)$ lies in a compact subset of $\mathcal{Q}$. So if $H.(X_c, q_c)$ is closed, then it is also compact and consquently $H.(X_c, q_c)$ is a periodic orbit. However, as the action of H preserve the marked vertical

saddle connections, the fact that $H.(X,q)$ is not periodic implies that $H.(X_c,q_c)$ is not periodic either. Therefore, $H.(X_c,q_c)$ is not closed. As the action of H preserve the marked vertical saddle connections, the H-orbit of (X,q) is not closed either. Consequently, the $SL(2,\mathbb{R})$-orbit of (X,q) is not closed in this case. This ends the proof.

□

6. Generic Veech groups

Just as flat surfaces of finite area, the typical flat surface with poles of higher order has a trivial Veech group. Theorem 2.4 provides a similar result to what Möller proved in [8]. As a result of our definition of a Veech group, we do not have to deal with hyperellipticity.

Proof of Theorem 2.4. First, we distinguish three kinds of chambers according to the shape of the core:
- chambers of the first kind, such that the core is not degenerated and its boundary is a unique closed saddle connection or a pair of parallel saddle connections.
- chambers of the second kind, such that the core is a unique saddle connection (in particular, the core is degenerated).
- chambers of the third kind otherwise.
We first deal with chambers of the second kind.

Chambers of the second kind are rare because $2g-2+n+p$ is a lower bound on the number of saddle connections of a surface of genus g with n conical singularities and p poles of higher order provided $n+p\geq 3$ (see Theorem 3.2). Surfaces with a unique saddle connection can only exist if $g=0$ and $n+p\leq 3$ and we assume in this paper that $n+p\geq 3$. Automorphisms of the sphere allow to fix three points so there is a unique surface up to scaling in strata such that $n+p=3$. All these surfaces have a unique saddle connection so their Veech group is of continuous type.

For a flat surface (X,q) that belongs to a chamber of the third kind, we look for two saddle connections of $\partial\mathcal{C}(X)$ whose holonomy vectors can be deformed independently from each other. Is should be recalled that saddle connections of $\partial\mathcal{C}(X)$ have no interior intersections.
Following Lemma 3.3 and Lemma 3.4, if the holonomy vectors of two saddle connections γ and γ' cannot be deformed independently from each other, then their union $\gamma\cup\gamma'$ cuts out X into several components one of which has trivial linear holonomy. The only case where all saddle connections of $\partial\mathcal{C}(X)$ are parallel is covered by chambers of the first kind (one unique saddle connection or two parallel saddle connections).
If the holonomy vectors of $\partial\mathcal{C}(X)$ are not dependent from each other (if there are at least two parameters of deformation), then a generic choice of periods will provide an irrational angle between the direction of a saddle connection γ and all the other saddle connections of $\partial\mathcal{C}(X)$ that are independant from γ. Since saddle connections that are not independant from γ share the same direction (Lemma 3.3), then such a configuration forbids a Veech group other than trivial. Therefore, the generic surface in a chamber of the third kind has a trivial Veech group.

We come back to chambers of the first kind. Since saddle connections of the boundary of the core share the same direction and the core is not degenerated, flat surfaces that belong to such chambers have a Veech group that is either trivial or of cyclic parabolic type (see Theorem 2.1).
Following Proposition 4.2, a flat surface (X,q) that belongs to such a chamber has a Veech group of cyclic parabolic type if and only if $core(X)$ splits into a finite number of cylinders

with commensurable moduli in the direction of the saddle connection of $\partial\mathcal{C}(X)$. If there are at least two cylinders in such a decomposition, we can broaden those cylinders in order to realize every ratio of moduli without leaving the chamber. In accordance with definitions of [8], we say that a flat surface (X, q) is generic in its stratum (or its chamber) if it lies outside a countable union of real codimension one submanifolds. There is a neighborhood of (X, q) in the stratum such that the locus of surfaces whose Veech group is of cyclic parabolic type is a countable union of real codimension one submanifolds (corresponding to commensurable moduli). Therefore, outside the locus of surfaces (X, q) where $core(X)$ is a unique cylinder, having a trivial Veech group is a generic property.

Let (X, q) be a surface such that $core(X)$ decomposes into a unique cylinder for the flow in the direction of the saddle connections of $\partial\mathcal{C}(X)$. Without loss of generality, we suppose that saddle connections of $\partial\mathcal{C}(X)$ are vertical. We should recall that the boundary of the cylinder does not necessarily coincide with the boundary of the core because there can be conical singularities in the interior of $core(X)$. If not all saddle connections of the boundary of the cylinder are parallel, then a generic perturbation of periods will produce a surface such that the decomposition into invariant components in the vertical direction is no longer merely a cylinder. Thus, in the neighborhood of such a flat surface (X, q) in the stratum, having a trivial Veech group is a generic property.

From now, we consider flat surfaces (X, q) such that:
- $\partial\mathcal{C}(X)$ is a unique saddle connection or a pair of parallel saddle connections.
- $core(X)$ decomposes into a unique cylinder for the flow in the direction of the saddle connections of $\partial\mathcal{C}(X)$.
- all saddle connections of the boundary of the cylinder are parallel.

Outside this locus $\mathcal{L}$, the generic surface has a trivial Veech group.
We note that this locus is an open set in which the Veech group of every surface is of cyclic parabolic type (Proposition 4.2). Such loci exist only in bidimensional strata. Indeed, there are only two independant complex parameters (periods of the parallel saddle connections of the boundary of the cylinder and the parameter prescribing the volume and the twist of the cylinder).
A stratum $\mathcal{Q}(a_1, \ldots, a_n, -b_1, \ldots, -b_p)$ of flat surfaces of genus g is bidimensional if and only if $g = 0$ and $n + p = 4$ or $g = n = p = 1$. Besides, there are at most two saddle connections in $\partial\mathcal{C}(X)$ so we have $p \leq 2$.

In surfaces of genus zero, cutting along a waist curve of the cylinder splits the surface into two connected components. Waist curves have a topological degree equal to zero so the sum of the orders of the singularities of each connected component is -2. Thus, we have two types of strata of surfaces of genus zero:
- $\mathcal{Q}(a, b, -a-2, -b-2)$ with $a, b \geq 1$
- $\mathcal{Q}(a, -1^2, -a-2)$ with $a \geq 1$

We have to add to them strata $\mathcal{Q}(a, -a)$ with $a \geq 2$ (because $\mathcal{Q}(1, -1)$ is empty) to get all strata in which some surfaces belong to locus $\mathcal{L}$.

In strata $\mathcal{Q}(a, b, -a-2, -b-2)$ with $a, b \geq 1$, for surfaces that belong to locus $\mathcal{L}$, cutting along a waist curve of the cylinder separates the domain of the pole of order $a+2$ (whose boundary is a unique closed saddle connection passing through the conical singularity of order a) from the domain the pole of order $b+2$ (whose boundary is a unique closed saddle connection passing through the conical singularity of order b). This qualitative shape of the core defines chambers. Every surface that belongs to these chambers is also in $\mathcal{L}$.
In strata $\mathcal{Q}(a, -1^2, -a-2)$ with $a \geq 1$, for surfaces that belong to locus $\mathcal{L}$, cutting along a waist curve of the cylinder separates the domain of the pole of order $a+2$ (whose boundary is a unique closed saddle connection passing through the conical singularity of order a) from



FIGURE 4. A surface in $\mathcal{Q}(2,-2)$ whose Veech group is of cyclic parabolic type.

a saddle connection connecting the two simple poles. This qualitative shape of the core also defines chambers. Every surface that belongs to these chambers is also in $\mathcal{L}$.
In strata $\mathcal{Q}(a,-a)$, every surface that belongs to locus $\mathcal{L}$ belongs to chambers such that the core is not degenerated and its boundary is a unique closed saddle connection or a pair of parallel saddle connections (chambers of the first kind as stated above). For every flat surface (X,q) in such a chamber, if we decompose $core(X)$ in the direction of the saddle connections of $\partial\mathcal{C}(X)$, we get exactly one invariant component. Indeed, there is at least one invariant component because all saddle connections of the boundary share the same direction and $core(X)$ is not degenerated. There is at most one invariant component because of Theorem 3.5. This invariant component is necessarily a cylinder because if it were a minimal component, there would be an additional parameter other than periods of the parallel saddle connections of $\partial\mathcal{C}(X)$ and the parameter prescribing the volume and the twist of the invariant component. Therefore, surfaces of $\mathcal{Q}(a,-a)$ that belong to locus $\mathcal{L}$ coincide with chambers such that the core is not degenerated and their boundary is a unique closed saddle connection or a pair of parallel saddle connections. This ends the proof.

□

Example 6.1. There is a chamber $\mathcal{C}$ in $\mathcal{Q}(2,-2)$ such that every surface in $\mathcal{C}$ has a Veech group of cyclic parabolic type, see Figure 4.

REFERENCES

[1] C. Boissy. *Configurations of saddle connections on quadratic differentials on $\mathbb{CP}^1$ and on hyperelliptic Riemann surfaces.* Commentarii Mathematici Helvitici, Volume 84, 757-791, 2009.

[2] C. Boissy. *Connected components of the strata of the moduli space of meromorphic differentials.* Commentarii Mathematici Helvitici, Volume 90, Issue 2, 255-286, 2015.

[3] J. Ellenberg, D.B. McReynolds. *Arithmetic Veech sublattices of $SL(2,\mathbb{R})$.* Duke Mathematical Journal, Volume 161, Number 3, 415-429, 2012.

[4] F. Haiden, L. Katzarkov, M. Kontsevich. *Flat surfaces and stability structures.* Preprint, arXiv:1409.8611, 2015.

[5] P. Hubert, E. Lanneau. *Veech groups without parabolic elements.* Duke Mathematical Journal, Volume 133, Number 2, 335-346, 2003.

[6] P. Hubert, T. Schmidt. *An introduction to Veech Surfaces.* Handbook of Modern Dynamics, Volume 1B, 2006.

[7] P. Hubert, G. Schmithüsen. *Infinite translation surfaces with infinitely generated Veech groups.* Journal of Modern Dynamics, Volume 4, Number 4, 715-732, 2010.

[8] M. Möller. *Affine groups of flat surfaces.* Handbook of Teichmüller theory, 369-387, 2009.

[9] H. Masur, A. Zorich. *Multiple saddle connections on flat surfaces and the principal boundary of the moduli spaces of quadratic differentials.* Geometric and Functional Analysis, Volume 18, Issue 3, 919-987, 2008.

[10] G. Schmithüsen. *An algorithm for finding the Veech group of an origami*. Preprint, 2003.
[11] K. Strebel. *Quadratic Differentials*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Volume 5, 1984.
[12] G. Tahar. *Counting saddle connections in flat surfaces with poles of higher order*. Preprint, arXiv:1606.03705, 2016.
[13] F. Valdez. *Veech groups, irrational billards and stable abelian differentials*. Discrete and Continuous Dynamical Systems, Series A, Volume 32, 1055-1063, 2012.
[14] W. Veech. *Teichmüller curves in modular space, Eisenstein series, and an application to triangular billiards*. Inventiones Mathematicae, Volume 97, 553-583, 1989.
[15] A. Zorich. *Flat Surfaces*. Frontiers in Physics, Number Theory and Geometry, 439-586, 2006.

(Guillaume Tahar) Institut de Mathématiques de Jussieu - UMR CNRS 7586
E-mail address: guillaume.tahar@imj-prg.fr